



UWS Academic Portal

The effects of microplastic on freshwater *Hydra attenuata* morphology & feeding

Murphy, Fionn; Prades, L.; Ewins, Ciaran; Quinn, Brian

Published in:

MICRO 2016: Fate and Impact of Microplastics in Marine Ecosystems

DOI:

[10.1016/B978-0-12-812271-6.00077-6](https://doi.org/10.1016/B978-0-12-812271-6.00077-6)

Published: 01/12/2016

Document Version

Publisher's PDF, also known as Version of record

[Link to publication on the UWS Academic Portal](#)

Citation for published version (APA):

Murphy, F., Prades, L., Ewins, C., & Quinn, B. (2016). The effects of microplastic on freshwater *Hydra attenuata* morphology & feeding. In J. Baztan, B. Jorgensen, S. Pahl, R. Thompson, & J-P. Vanderlinden (Eds.), *MICRO 2016: Fate and Impact of Microplastics in Marine Ecosystems: From the Coastline to the Open Sea* (pp. 78-79). Elsevier B.V.. <https://doi.org/10.1016/B978-0-12-812271-6.00077-6>

General rights

Copyright and moral rights for the publications made accessible in the UWS Academic Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact pure@uws.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

The effects of microplastic on freshwater *Hydra attenuata* feeding & morphology

Fionn Murphy¹, Laura Prades¹, Ciaran Ewins¹, and Brian
Quinn¹

¹ Institute of Biomedical and Environmental Health Research, University of the West of Scotland,
Paisley, PA1 2BE, Scotland

Contact: fionn.g.murphy@gmail.com

Microplastic in the Environment:

Marine

- Accumulating on shorelines worldwide
- Many marine organisms ingesting microplastic
- Most research focused here

Freshwater

- Present in lakes and rivers
- Research lacking

Knowledge Gap

- There are currently no standardised tests for assessing the impact of microplastic

Current Research:

- **Question:**
 - Can microplastic have an impact on freshwater organisms?
- Investigating the effect of microplastics on the feeding & morphology of a freshwater cnidarian, *Hydra attenuata*
- **Aims:**
 - Determine if *Hydra* can ingest microplastic
 - Determine if microplastic can effect the feeding of *Hydra*
 - Determine if microplastic can effect the morphology of *Hydra*

Hydra attenuata:

- Used widely in bioassays to test the toxicity of environmental contaminants (Quinn et al., 2008)



Hydra Feeding & Morphology:

- Fed *Artemia salina*




Hydra morphology – Toxicity to polyp phase

| | | | | | | | | | | |
|---------------------------------------|--------------------------------------|--|----------------------|------------------------------|---------------------------------------|--|------------------------------|---------------------------------|-----------------|---------------|
| | | | | | | | | | | |
| Normal | Increasing Degree of Toxicity | | | | | | Osmoregulation loss | Terminal Stages | | |
| Extended tentacles and body reactive. | Partially contracted slow reactions. | Clubbed tentacles. Body slightly contracted. | Shortened tentacles. | Tentacles and body shortened | Totally contracted tentacles visible. | Totally contracted no visible tentacles. | Expanded, tentacles visible. | Expanded, no visible tentacles. | Dead but intact | Disintegrated |
| Score 10 | Score 9 | Score 8 | Score 7 | Score 6 | Score 5 | Score 4 | Score 3 | Score 2 | Score 1 | Score 0 |

Wilby (1988)

Microplastic (MP):


- MP <400 μm




Exposure:

- *Hydra* starved for three days
- Placed in 0.5 ml Eppendorf tubes containing MP
- Concentrations: Control, 0.01, 0.02, 0.04, 0.08 g/ml
- 10 *Artemia* added
- Ingestion of MP & *Artemia* checked after 30 & 60 min
- Morphology recorded after 3, 24, 48, & 96 hrs


Feeding Rate:



Ingestion of MP:




Morphology Scores:




Other observations:

- Buoyancy change
- Attachment to microplastic



Conclusions:

- *Hydra* are capable of ingesting microplastic
- The presence of microplastic disrupted the feeding of the *Hydra*



Implications for the environment:

- MP can potentially have an effect on feeding
- Feeding is an ecologically relevant endpoint
- Could alter community structure

Thank You

Questions?

Email: fionn.g.murphy@gmail.com

Acknowledgements

St-Lawrence Centre (SCL) of Montreal, Quebec for providing the Hydra used in this study

References

QUINN, B., GAGNÉ, F. & BLAISE, C. 2008. An investigation into the acute and chronic toxicity of eleven pharmaceuticals (and their solvents) found in wastewater effluent on the cnidarian, *Hydra attenuata*. *Science of the total environment*, 389, 306-314.

Wilby, O. K. "The Hydra regeneration assay." *Proceedings of workshop organised by Association Francaise de Teratologie*. 1988.